



once-daily

NUZYRA[®]

(omadacycline) 100 mg for injection
150 mg tablets

PATIENT GUIDE

Starting treatment with NUZYRA

What is NUZYRA?

NUZYRA[®] (omadacycline) is a prescription antibiotic medicine used to treat adults 18 years and older with:

- An infection of the lung caused by certain kinds of germs called bacteria (community-acquired bacterial pneumonia)
- Skin infections caused by certain bacteria (acute bacterial skin and skin structure infections)

Some infections are caused by viruses and not by bacteria. Examples include viral infections such as the common cold or flu. Antibacterial medicines, including NUZYRA, do not treat viral infections. NUZYRA should only be used to treat bacterial infections. Call your healthcare provider if you think your condition is not getting better while you are taking NUZYRA.

Please see Important Safety Information on pages 8-9.

INSIDE YOU'LL FIND THE FOLLOWING INFORMATION:

What is NUZYRA® (omadacycline)?.....	3
Taking NUZYRA Tablets.....	4
Information About Taking NUZYRA.....	6
Important Safety Information for NUZYRA.....	8
Next Steps—How to Get NUZYRA	10

WHO SHOULD NOT TAKE NUZYRA?

Do not take NUZYRA if you are allergic to tetracycline-class drugs. Ask your doctor or pharmacist for a list of these medicines if you are not sure.

Please see Important Safety Information on pages 8-9.



WHAT IS NUZYRA® (omadacycline)?

NUZYRA is an antibiotic used to treat adults who have one of the following conditions:



COMMUNITY-ACQUIRED BACTERIAL PNEUMONIA (CABP)

A type of lung infection that causes inflammation of the lungs

- Symptoms can include:**
- Chest pain
 - Cough
 - Fever
 - Difficulty breathing



ACUTE BACTERIAL SKIN AND SKIN STRUCTURE INFECTIONS (ABSSSI)

A type of skin infection

- Symptoms at the infection site can include:**
- Pain/tenderness
 - Redness
 - Inflammation

NUZYRA works by stopping the growth of certain bacteria that commonly cause these infections.

Antibiotic medicines like NUZYRA should only be used to treat bacterial infections. They do not treat viral infections such as the common cold.

TAKING NUZYRA TABLETS



HOW DO I TAKE NUZYRA?

Take **NUZYRA® (omadacycline)** with water only. You should take **NUZYRA on an empty stomach.**

- Do not eat or drink for at least 4 hours before and 2 hours after taking NUZYRA (except water)
- Do not eat or drink any dairy products, antacids, or multivitamins for 4 hours after taking NUZYRA tablets
- Be sure to talk to your doctor if you miss a dose



WHEN DO I TAKE NUZYRA?

Because you cannot eat or drink (except water) for at least 4 hours before—and 2 hours after—taking NUZYRA, the time you take this medication depends on your lifestyle. You may find it more convenient to take NUZYRA in the morning or at bedtime. Talk to your healthcare provider or pharmacist to decide what time works best for you.

REMEMBER

- Keep NUZYRA tablets stored at 68 °F to 77 °F, and do not freeze
- No food or drink (except water) for at least 4 hours, and then take with water
- NUZYRA can be taken at bedtime or upon waking
- No food or drink (except water) for 2 hours after taking NUZYRA
- No dairy products, antacids, or multivitamins for 4 hours after dosing



Please see *Important Safety Information on pages 8-9.*

YOUR DAILY TREATMENT LOG

Keep track of your treatment by checking off each day's dose.

Although it's common to feel better early on in the course of therapy, don't skip a dose, and be sure to finish all of your medication.

COMMUNITY-ACQUIRED BACTERIAL PNEUMONIA (CABP)

2 tablets **TWICE** on Day 1

- DOSE 1 
- DOSE 2 

Then 2 tablets together each day for remainder of treatment

- DAY 2 
- DAY 3 DAY 4
- DAY 5 DAY 6
- DAY 7 DAY 8
- DAY 9 DAY 10
- DAY 11 DAY 12
- DAY 13 DAY 14

ACUTE BACTERIAL SKIN AND SKIN STRUCTURE INFECTIONS (ABSSSI)

3 tablets on Day 1 & Day 2

- DAY 1 
- DAY 2 

Then 2 tablets together each day for remainder of treatment

- 
- DAY 3 DAY 4
- DAY 5 DAY 6
- DAY 7 DAY 8
- DAY 9 DAY 10
- DAY 11 DAY 12
- DAY 13 DAY 14

Tablets are not actual size.

NUZYRA TREATMENT DURATION IS TYPICALLY 7-14 DAYS; PLEASE TAKE NUZYRA EXACTLY AS DIRECTED BY YOUR DOCTOR.

INFORMATION ABOUT TAKING NUZYRA

Please be aware of the following information while taking NUZYRA:



Common Side Effects

The most common side effects of NUZYRA include: nausea, vomiting, increased liver tests, elevated blood pressure (hypertension), headache, diarrhea, difficulty sleeping (insomnia), and constipation.

- Nausea and vomiting are more common in patients who receive the oral loading doses (Days 1 and 2)
- Reactions at the infusion site (pain, swelling, redness, and irritation) were also common in patients who received NUZYRA intravenously (IV)
- These are not all of the possible side effects of NUZYRA. Tell your healthcare provider about any side effect that bothers you or that does not go away



Possible Allergic Reactions to Tetracyclines

NUZYRA belongs to the tetracycline family of antibiotics, and may cause similar adverse reactions. Be sure to tell your doctor if you have ever had an allergic reaction to an antibiotic known as a tetracycline.

- For example, doxycycline and minocycline are some common tetracyclines



Other Medicine

Be sure to tell your doctor about the medicines you take, including prescription drugs (such as oral contraceptives) and over-the-counter medicines, vitamins, and herbal supplements. It is especially important to mention if you are taking any of the following:

- Blood thinners (anticoagulants) such as warfarin
- Antacids or mineral supplements



Breastfeeding and Pregnancy

Do not breastfeed while taking NUZYRA and for 4 days after the last dose.

If you are pregnant or plan to become pregnant, NUZYRA may harm an unborn baby.



Diarrhea

Tell your doctor if you have diarrhea or have had diarrhea while taking antibiotics. Be sure to mention if you experience any of the following, even after you have stopped taking NUZYRA:

- Diarrhea
- Stomach cramps
- Bloody stools



Antibacterial Resistance

Remember that although it is common to feel better after only a few doses of your antibiotic, NUZYRA should be taken exactly as directed.

- Skipping doses or not finishing all of the medication prescribed may:
 - Make the treatment not work as well
 - Increase the likelihood that bacteria will develop resistance and will not be treatable by NUZYRA or other antibacterial drugs in the future

Please see Important Safety Information on pages 8-9.



IMPORTANT SAFETY INFORMATION FOR NUZYRA® (omadacycline)

What is NUZYRA?

NUZYRA® (omadacycline) is a prescription antibiotic medicine used to treat adults 18 years and older with:

- An infection of the lung caused by certain kinds of germs called bacteria (community-acquired bacterial pneumonia)
- Skin infections caused by certain bacteria (acute bacterial skin and skin structure infections)

Some infections are caused by viruses and not by bacteria. Examples include viral infections such as the common cold or flu. Antibacterial medicines, including NUZYRA, do not treat viral infections. NUZYRA should only be used to treat bacterial infections. Call your healthcare provider if you think your condition is not getting better while you are taking NUZYRA.

Who should not take NUZYRA?

Do not take NUZYRA if you are allergic to tetracycline class drugs. Ask your doctor or pharmacist for a list of these medicines if you are not sure.

What is the most important information I should know about NUZYRA?

In the community-acquired bacterial pneumonia clinical trial, death occurred in 2% of NUZYRA-treated patients (8 out of 382) compared to 1% of patients (4 out of 388) who received another antibiotic. All deaths in both treatment groups occurred in patients older than 65 years of age.

What should I tell my healthcare provider before taking NUZYRA?

Before taking NUZYRA, tell your healthcare provider:

- **About all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as NUZYRA may harm an unborn baby. Taking NUZYRA during tooth development (during the last half of pregnancy, infancy and during childhood up to the age of 8 years) may cause serious side effects including permanent discoloration of teeth and slow bone growth of your baby. Talk to your doctor before taking NUZYRA if you plan to become pregnant, or if you are already taking NUZYRA and plan to become pregnant.
 - are breast feeding or plan to breast feed. NUZYRA passes into your milk and may harm your baby. You and your doctor should decide if you will take NUZYRA or breastfeed. You should not do both.
 - have diarrhea or have had diarrhea while taking antibiotics, as you can develop an infection of the intestines.

- **About all the medicines you take**, including prescription drugs (such as oral contraceptives) and over-the-counter medicines, vitamins, and herbal supplements. NUZYRA and certain other medicines may affect each other or may cause side effects. **It is especially important to mention** if you are taking any of the following:

- blood thinners (anticoagulants) such as warfarin
- antacids that contain aluminum, calcium, or magnesium or iron-containing products.

What are the possible side effects of NUZYRA?

NUZYRA may cause serious side effects including:

- **Serious allergic reactions** have occurred in patients taking NUZYRA and life-threatening allergic reactions have occurred with other tetracyclines. Stop taking NUZYRA and get emergency medical help if a serious allergic reaction occurs.
- **Clostridioides difficile-associated diarrhea (CDAD)** - CDAD is an infection of your intestines (bowels) that can happen with many antibacterial medicines like NUZYRA and may cause mild diarrhea to life-threatening swelling of your intestines (colitis). Call your healthcare provider right away if you get stomach cramps, fever, watery diarrhea, diarrhea that does not go away, or bloody stools.
- NUZYRA is similar to other tetracycline class antibacterial drugs and may cause similar side effects. NUZYRA can make your skin sensitive to the sun and the light from sunlamps and tanning beds. Protect your skin while out in sunlight. Stop taking NUZYRA and tell your healthcare provider right away if you have blurred vision, vision loss, or unusual headaches or if you get stomach pain, yellowing of your skin or eyes, unexplained bleeding, confusion, or loss of appetite, or a flat red skin patch that recurs at the same site or sites each time the medication is restarted.

The most common side effects of NUZYRA include: nausea, vomiting, increased liver tests, elevated blood pressure (hypertension), headache, diarrhea, difficulty sleeping (insomnia), and constipation. Reactions at the infusion site (pain, swelling, redness and irritation) were also common in patients who received NUZYRA intravenously (IV).

These are not all of the possible side effects of NUZYRA.

Call your healthcare provider if you have these or any other side effects while taking NUZYRA.

You are encouraged to report negative side effects of NUZYRA. Please contact Paratek Pharmaceuticals at 1-833-727-2835 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please see accompanying full Prescribing Information in pocket.



once-daily
NUZYRA®
(omadacycline)
100 mg for injection / 150 mg tablets



once-daily

NUZYRA®

(omadacycline) 100 mg for injection
150 mg tablets

NEXT STEPS



Your healthcare professional has sent your prescription to a network pharmacy.

The pharmacy may call you to review your insurance coverage.



It's important to speak with the pharmacy promptly to ensure that all of the necessary information is available to fill your NUZYRA prescription quickly.

If you haven't heard from the pharmacy within 24 hours, call **NUZYRA Central® Support Services** at **1-877-4-NUZYRA (1-877-468-9972)**.



NUZYRA may be delivered directly to your home at no additional cost.

COPAY ASSISTANCE MAY BE AVAILABLE FOR ELIGIBLE PATIENTS

Commercially insured patients may pay as little as [\$0].*

*Terms and conditions apply.

Please see Important Safety Information on pages 8-9.



© 2025 Paratek Pharmaceuticals, Inc. All rights reserved.
PARATEK® and the hexagon logo are registered trademarks of Paratek Pharmaceuticals, Inc.
NUZYRA® and its design logo are registered trademarks of Paratek Pharmaceuticals, Inc.

US-NUA-0589-01 05/25